

FIGURE 1 – Plan map of the ACKIO mineralized surface expressions




TABLE 1 – Drill collar details, composite radioactivity and U₃O₈ assay results, drill hole AK22-069

DDH	Target Area	East	North	Elevation	Azimuth	Dip	EOH	Radioactivity (>300 cps)	Assay Results (>0.05 wt% U ₃ O ₈)
AK22-69	ACKIO	526,009	6,372,980	465	90	-50	327	No significant results	0.08% over 1.0 m at 56.0 m
								643 cps over 1.5 m at 81.4 m	0.09% over 1.7 m at 81.3 m
								550 cps over 0.1 m at 87.0 m	0.09% over 1.0 m at 86.5 m
								1,687 cps over 26.9 m at 90.9 m¹	0.90% over 31.0 m at 90.5 m¹
								includes 2,614 cps over 0.45 m at 96.0 m	0.86% over 0.5 m at 96.0 m
								and includes 3,100 cps over 11.3 m at 104.85 m²	1.86% over 12.5 m at 104.0 m²
								550 cps over 0.4 m at 121.1 m	No significant results
								653 cps over 0.2 m at 186.7 m	Assay results pending
								550 cps over 0.15 m at 200.15 m	Assay results pending
								950 cps over 0.15 m at 209.65 m	Assay results pending
1							327.0	1 DDH	

NOTES: East and North units are metres using NAD83 datum, UTM Zone 13N

Elevation is recorded as "metres above sea level"

EOH = End of hole, measured in metres

Composite radioactivity results use 300 cps cut-off and do not contain greater than 2.0 m consecutive dilution (i.e., dilution is <300 cps)

Composite U₃O₈ results use 0.05% U₃O₈ cut-off and do not contain greater than 2.0 m consecutive dilution (i.e., dilution is <0.05% U₃O₈)

"includes/and" are composite U₃O₈ results using 0.50% U₃O₈ cut-off and do not contain greater than 2.0 m consecutive dilution (i.e., dilution is <0.50% U₃O₈)

¹includes 7.1 m core loss over interval length

²includes 2.25 m core loss over interval length